arXiv:1210.3216v2 [quant-ph] 13 Oct 2012

Is composite noise necessary for sudden death of entanglement?

Yashodamma K.O[1] and Sudha[1, 2, *]
[1]Department of Physics, Kuvempu University, Shankaraghatta, Shimoga-577 451, India.
[2]Inspire Institute Inc., Alexandria, Virginia, 22303, USA.

(Dated: October 16, 2012)

It is shown that finite time disentanglement or entanglement sudden death can occur even when only one part of the composite entangled state is exposed to a noisy environment. The result is illustrated through the action of different noisy environments individually on a single qubit of the so-called X class of states. Composite noise on a part or whole of the entangled state is thus not found to be necessary for sudden death of entanglement.

PACS numbers: 03.67.Yz, 03.65.Ud,42.50.Lc

I. INTRODUCTION

Quantum entanglement [1, 2], a special kind of correlation that exists in composite quantum systems has been of great relevance both for its importance in foundational issues of quantum mechanics [1, 2] and for its use as resource in quantum computation [3, 4]. The fragility of entangled quantum systems [5, 6] to the environment surrounding them is the main cause of concern for realizing their technological aspirations. Decoherence refers to the process through which superposition of quantum states are irreversibly transformed into statistical mixtures due to coupling of quantum system with surrounding environment [5–9]. Eventhough it is not possible to fully characterize the environment, different models have been proposed. Early stage disentanglement or Entanglement Sudden Death (ESD) [10, 11] is a consequence of the exposure of entangled systems to noisy environment or channels.

In [10], Yu and Eberly elucidated the concept of finite time disentanglement by examining how two entangled qubits individually interacting with pure vacuum noise lose their entanglement in a finite time. Further examination of ESD has revealed that this early stage disentanglement is an illustration of nonadditivity of individual coherence decay rates [11]. This inevitable loss of entanglement has gained lot of attention in the quantum information community and efforts are being made to find ways in which ESD can be avoided [12].

It is to be noted here that the studies on finite time disentanglement in bipartite systems [10, 11, 13–20] concentrated on situations in which either both the subsystems are coupled individually to one or more noisy environments or one of the subsystems is coupled to more than one noise. While it can be interpreted that nonadditivity of individual coherence decay rates is manifested in the occurence of ESD in such cases [11], a question of interest is whether exponential decay of entanglement always results when one of the subsystems alone is coupled to a noisy environment leaving the other subsystem entirely free of noise. Except in Ref. [21, 22], where sudden death of entanglement in qubit-qutrit states when only one of them is either subjected to dephasing [21] or depolarizing noise [22], not much attention is paid to the finite time disentanglement due to a single local noise. We wish to take up this issue and examine the disentanglement dynamics of bipartite states when one of the subsystems is kept free of noise and the other is subjected to a *single* noisy environment. We have shown that sudden death of entanglement results even when one part of a composite entangled system is exposed to a single quantum noise.

The article is organized as under: Sec.I contains introductory remarks. In Sec.II we have examined the effect of amplitude, phase and depolarizng noises each acting locally on a qubit of two-qubit X states. Two special states of this class, the isotropic, Werner states are analyzed under the local action of these three noises and their disentanglement times are tabulated in Sec.III. Concluding remarks are given in Sec.IV.

II. SUDDEN DEATH OF ENTANGLEMENT IN A CLASS OF X STATES

We have considered the class of two qubit states of the form,

$$\rho = \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & b & z & 0 \\ 0 & z^* & c & 0 \\ 0 & 0 & 0 & d \end{bmatrix} \tag{1}$$

where $a+b+c+d=1$, $z=x+iy$. This is a subset of the so-called X class of states [10, 11, 16] and it encompasses both pure states such as the all important Bell states and mixed states such as Werner and isotropic states.

It is readily seen that concurence [23, 24], a well known measure of entanglement for two-qubit states, of the above state is of the form

$$C = 2\,\mathrm{Max}\left[0, |z| - \sqrt{ad}\right]. \tag{2}$$

The noisy channels we have considered here are amplitude noise, phase noise and depolarizing noise. It is well

---

[*] Electronic address: arss@rediffmail.com

known that the effect of each of these noises on a quantum system can be characterized through the corresponding Kraus operators [3, 10**?** , 11].

A. Amplitude Noise:

The process such as spontaneous emission of a photon is characterized by the quantum operation so called amplitude damping or amplitude noise. It gives the right description for energy dissipation from a quantum system. The Kraus operators for a single qubit amplitude noise are given by [10]

$$E_0 = \begin{bmatrix}\eta & 0\\ 0 & 1\end{bmatrix}; E_1 = \begin{bmatrix}0 & 0\\ \sqrt{1-\eta^2} & 0\end{bmatrix} \tag{3}$$

where $\eta = e^{-\frac{\tau t}{2}}$ and τ denotes the longitudinal decay rate.

When only the first qubit is subjected to amplitude noise leaving the second qubit noise free, the corresponding Kraus operators are given by

$$\begin{aligned} K_{1a} &= \begin{bmatrix}\eta & 0\\ 0 & 1\end{bmatrix} \otimes \begin{bmatrix}1 & 0\\ 0 & 1\end{bmatrix} \\ K_{2a} &= \begin{bmatrix}0 & 0\\ \sqrt{1-\eta^2} & 0\end{bmatrix} \otimes \begin{bmatrix}1 & 0\\ 0 & 1\end{bmatrix} \end{aligned}$$

The time evolved density matrix with amplitude noise acting on first qubit of state Eq. (1) is obtained as,

$$\begin{aligned} \rho_a(t) &= K_{1a}\rho K_{1a}^\dagger + K_{2a}\rho K_{2a}^\dagger \\ &= \begin{bmatrix} a\eta^2 & 0 & 0 & 0 \\ 0 & b\eta^2 & z\eta & 0 \\ 0 & z^*\eta & c+a-a\eta^2 & 0 \\ 0 & 0 & 0 & d+b-b\eta^2 \end{bmatrix} \end{aligned}$$

It can be readily seen that the state $\rho_A(t)$ retains its X-form and hence its concurrence can be calculated using Eq. (2). We get,

$$C_a = 2\,\mathrm{Max}\left[0, \eta\left(|z| - \sqrt{a[b+d-b\eta^2]}\right)\right]. \tag{4}$$

It can be seen that $C_a = 0$ when $|z| - \sqrt{a(b+d-b\eta^2)} \leq 0$ or $t \geq \frac{1}{\tau}\ln\left[\frac{ab}{a(b+d)-|z|^2}\right]$. Thus, depending on the parameters of the initial density matrix, one can either observe sudden death of entanglement or not. This feature is illustrated in Fig. 1.



FIG. 1: When the density matrix parameters are chosen to be $a = 0.1, b = 0.4, c = 0.4, d = 0.1$ and $|z|^2 = 0.04$ (solid line) the entanglement of the state vanishes when $\Gamma t = 1.386$. But when the density matrix parameters are chosen to be $a = 0.1,\ b = 0.2,\ c = 0.6,\ d = 0.1$ and $|z|^2 = 0.04$ there is an exponential decay of entanglement (dashed line) contrary to the previous case.

B. Phase Noise

Phase damping or phase noise is a uniquely quantum mechanical noise that describes the loss of quantum information without loss of energy. Under the action of phase noise, the off-diagonal elements of the density matrix decay exponentially with time. The Kraus operators for single qubit phase noise are given by [**?**]

$$K_0 = \begin{bmatrix}1 & 0\\ 0 & \gamma\end{bmatrix}; \quad K_1 = \begin{bmatrix}0 & 0\\ 0 & \sqrt{1-\gamma^2}\end{bmatrix} \tag{5}$$

where $\gamma = e^{-\frac{\tau t}{2}}$ and τ denotes transvere decay rate.

The Kraus operators for the action of phase noise on the first qubit alone being $K_{1p} = K_1 \otimes I_2$, $K_{2p} = K_1 \otimes I_2$ (I_2 denotes the 2×2 identity matrix), we have

$$\begin{aligned} \rho_p(t) &= K_{1p}\rho K_{1p}^\dagger + K_{2p}\rho K_{2p}^\dagger \\ &= \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & b & z\gamma & 0 \\ 0 & z^*\gamma & c & 0 \\ 0 & 0 & 0 & d \end{bmatrix} \end{aligned} \tag{6}$$

We can observe that only the off diagonal terms are affected but the time evolved density matrix retains its X-form. On evaluating the concurrence for this state, we get

$$C_p = 2\,\mathrm{Max}\left[0, \gamma|z| - \sqrt{ad}\right] \tag{7}$$

Notice that $C_P = 0$ when $\left(\gamma|z| - \sqrt{ad}\right) \leq 0$ or $t \geq \frac{1}{\tau}\ln\left[\frac{|z|^2}{ad}\right]$. As in the case of amplitude noise, depending on the initial parameters of the density matrix, one can either observe sudden dealth of entanglement or not.

The illustration of the result is done for some specified values of $a,\ b,\ c,\ d$ and $|z|^2$ in the following plots.

C. The Depolarizing noise

A quantum noise that converts a qubit into a completely mixed state with probability p leaves it untouched



FIG. 2: While the density matrix parameters $a = 0.2, b = 0.3, c = 0.3, d = 0.2$ and $|z|^2 = 0.09$ give rise to sudden death of entanglement (solid line), there is no such finite time disentanglement in the density matrix ρ with $a = 0.5, b = 0.1, c = 0.4, d = 0$ and $|z|^2 = 0.01$

with probability $1 - p$ is the depolarizing noise [3]. The Kraus operators for depolarizing noise are given by [3]

$$\begin{aligned} D_1 &= \sqrt{1-p}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \quad D_2 = \sqrt{\frac{p}{3}}\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \\ D_3 &= \sqrt{\frac{p}{3}}\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}; \quad D_4 = \sqrt{\frac{p}{3}}\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \qquad (8) \end{aligned}$$

To observe the effect of depolarizing noise on entanglement when it acts only on the first qubit of a composite system, we have to consider the Kraus operators,

$$\begin{aligned} K_{1d} &= D_1 \otimes I_2, \quad K_{2d} = D_2 \otimes I_2, \\ K_{3d} &= D_3 \otimes I_2, \quad K_{2d} = D_4 \otimes I_2. \end{aligned}$$

The time evolved density matrix with depolarizing noise acting on first qubit alone is given by,

$$\rho_d(t) = \sum_{i=1}^{4} K_{id}\, \rho\, K_{id}^{\dagger} \qquad (9)$$

$$= \begin{pmatrix} a + \frac{2p(c-a)}{3} & 0 & 0 & 0 \\ 0 & b + \frac{2p(d-b)}{3} & \frac{(3-4p)z}{3} & 0 \\ 0 & \frac{(3-4p)z^*}{3} & c + \frac{2p(a-c)}{3} & 0 \\ 0 & 0 & 0 & d + \frac{2p(b-d)}{3} \end{pmatrix}$$

As the resultant density matrix has retained its X-form even under the depolarizing noise, we can readily obtain the concurrence of the state. It is given by

$$c_d = \frac{2}{3}\,\mathrm{Max}\left[0,\ (3-4p)|z| - \sqrt{[3a + 2p(c-a)][3d + 2p(b-d)]}\right] \qquad (10)$$

One can notice that sudden death of entanglement is a quite possible happening in this case. We have illustrated two situations where initial entanglement decays in finite time due to depolarizing noise.

The family of X states considered in Eq.(1) include quantum states of importance such as Bell states, Werner states and isotropic states. In what follows, we will show that each of these states exhibit sudden death of entanglement under the action of a single noise on a single qubit.



FIG. 3: While the initial density matrix parameters are chosen to be $a = 0.1, b = 0.6, c = 0.2, d = 0.1$ and $|z|^2 = 0.04$ in one case(solid line), they are $a = 0, b = 0.5, c = 0.5, d = 0$ and $|z|^2 = 0.2$ in the other (dot-dashed line). Sudden death of entanglement is observed in both the cases.

III. ISOTROPIC AND WERNER STATES

Isotropic State: The isotropic states are d^2 dimensional bipartite states that are convex mixtures of a maximally entangled state with a maximally mixed state [25]. They are conveniently expressed in the form [26]

$$\rho_{\mathrm{iso}} = \frac{1-x}{d^2-1}\left[I - |\psi\rangle\langle\psi|\right] + x|\psi\rangle\langle\psi| \qquad (11)$$

where $0 \leq x \leq 1$ and $|\psi\rangle$ is any maximally entangled state of dimension d^2. Notice that ρ_{iso} is entangled when the parameter x is greater than $\frac{1}{d}$. Isotropic states possess the property of invariance under unitary transformations of the form $U \otimes U^*$.

A two-qubit isotropic state is given by

$$\rho = \begin{pmatrix} \frac{1-x}{3} & 0 & 0 & 0 \\ 0 & \frac{2x+1}{6} & \frac{4x-1}{6} & 0 \\ 0 & \frac{4x-1}{6} & \frac{2x+1}{6} & 0 \\ 0 & 0 & 0 & \frac{1-x}{3} \end{pmatrix}. \qquad (12)$$

The concurrence of the above state is readily seen to be $C_i = \mathrm{Max}\,[0,\, 2x - 1]$ which implies $C_i > 0$ for all $x > \frac{1}{2}$. An interaction of the state's first qubit alone with amplitude noise leads to

$$\rho_{ia} = \begin{pmatrix} \frac{(1-x)\eta^2}{3} & 0 & 0 & 0 \\ 0 & \frac{(2x+1)\eta^2}{6} & \frac{(4x-1)\eta}{6} & 0 \\ 0 & \frac{(4x-1)\eta}{6} & \frac{(3+2(x-1))\eta^2}{6} & 0 \\ 0 & 0 & 0 & \frac{3-(1+2x)\eta^2}{6} \end{pmatrix}. \qquad (13)$$

The concurrence of ρ_{ia} is given by

$$C_{ia} = \frac{\eta}{3}\,\mathrm{Max}\,\left[0,\ (4x-1) - \sqrt{2(1-x)(3-(1+2x)\eta^2)}\right] \qquad (14)$$

Notice that $C_{ia} = 0$ when

$$(4x-1) - \sqrt{2(1-x)(3-(1+2x)\eta^2)} \leq 0$$

or when

$$t \geq \frac{1}{\tau}\ln\left[\frac{2(2x+1)(x-1)}{6(x-1) + (4x-1)^2}\right]. \qquad (15)$$

It is not difficult to observe that for all values $x < 0.625$, sudden death of entanglement occurs while for $x \geq 0.625$, there is an exponential decay of entanglement. This result is illustrated in the following graph.



FIG. 4: Variation of entanglement with respect to time when the first qubit of the isotropic state is subjected to amplitude noise.

Now subjecting the first qubit of the isotropic state to phase noise we have,

$$\rho_{ip} = \frac{1}{6}\begin{pmatrix} 2(1-x) & 0 & 0 & 0 \\ 0 & 2x+1 & (4x-1)\gamma & 0 \\ 0 & (4x-1)\gamma & 2x+1 & 0 \\ 0 & 0 & 0 & 2(1-x) \end{pmatrix}. \tag{16}$$

The concurrence of this state is readily seen to be,

$$C_{ip} = \frac{1}{3}\,\mathrm{Max}\,[0, (4x-1)\gamma - 2(1-x)]\,, \tag{17}$$

and it is clear that, $C_{ip} = 0$ when $(4x-1)\gamma - 2(1-x) \leq 0$ or when $t \geq \frac{2}{\tau}\ln\left[\frac{4x-1}{2-2x}\right]$. One can thus calculate the time at which entanglement vanishes under the action of phase noise on the first qubit. Except at $x = 1$, where we have an exponential decay of entanglement, one can see a finite time disentanglement for all values of x in the open interval $(0.5, 1)$. The following graph illustrates this result.



FIG. 5: Variation of entanglement with respect to time when the first qubit of the isotropic state is subjected to phase noise. There is strictly exponential decay of entanglement only when $x = 1$. For all other values of x, there is finite time disentanglement.

When we allow the first qubit of the isotropic state to interact with depolarizing noise, the time evolved density matrix $\rho_{id} = \sum_{i=1}^{4} K_{id}\,\rho\, K_{id}^{\dagger}$ retains its X form (see Eq.(1)) with

$$\begin{aligned} a &= d = \frac{3-3x+p(4x-1)}{9}; z = \frac{(3-4p)(4x-1)}{18} \\ b &= c = \frac{3+6x+2p(1-4x)}{18}; \end{aligned} \tag{18}$$

A simplified expression for concurrence is given by

$$C_{id} = \frac{1}{3}\,\mathrm{Max}\,\left[0,\, 2p(1-4x) + 6x - 3\right]. \tag{19}$$

With $p = 1 - e^{-\tau\, t}$, it can be seen that $c_{id} = 0$ when $t \geq \frac{1}{\tau}\ln\left(\frac{8x-2}{2x+1}\right)$. It can also be seen that sudden death of entanglement occurs for the state for all values of $x > \frac{1}{2}$. We have illustrated it through the following graph.



FIG. 6: Depolarizing noise even on a single qubit of the isotropic state leads to sudden death of entanglement for all values of the parameter $\frac{1}{2} < x \leq 1$.

Werner State: Werner states are d^2 dimensional bipartite states that remain invariant under unitary transformations of the form $U \times U$ [27]. A two-qubit Werner state is a mixture of the fully mixed state $\frac{I}{4}$ with probability $(1-x)$ and a singlet state $|\psi\rangle = \frac{|01\rangle - |10\rangle}{\sqrt{2}}$ with probability x [28].

$$\rho_w = (1-x)\frac{I}{4} + x|\psi\rangle\langle\psi| \tag{20}$$

Explicitly, ρ_w is given by

$$\rho_w = \frac{1}{4}\begin{pmatrix} 1-x & 0 & 0 & 0 \\ 0 & 1+x & 2x & 0 \\ 0 & 2x & 1+x & 0 \\ 0 & 0 & 0 & 1-x \end{pmatrix} \tag{21}$$

As $C_w = \mathrm{Max}\,[0, \frac{3x-1}{2}]$ we have $C_w \neq 0$ for all $x > \frac{1}{3}$.

On allowing the first qubit of the Werner state to interact with amplitude noise we have

$$\rho_{wa} = \begin{pmatrix} \frac{(1-x)\eta^2}{4} & 0 & 0 & 0 \\ 0 & \frac{(1+x)\eta^2}{4} & \frac{x\eta}{2} & 0 \\ 0 & \frac{x\eta}{2} & \frac{2+(x-1)\eta^2}{4} & 0 \\ 0 & 0 & 0 & \frac{2-(1+x)\eta^2}{4} \end{pmatrix}, \tag{22}$$

and

$$C_{wa} = \frac{\eta}{2}\,\mathrm{Max}\left[0,\, 2x - \sqrt{(1-x)(2-(1+x)\eta^2)}\right] \tag{23}$$

. Notice that $C_{wa} = 0$ when

$$2x - \sqrt{(1-x)(2-(1+x)\eta^2)} \leq 0$$

or when

$$t \geq \frac{1}{\tau} \ln \left[\frac{1-x^2}{2-2x-4x^2} \right].$$

The range of values of x for which finite time disentanglement occurs in this case is shown in Fig. 7.



FIG. 7: The action of amplitude noise on a single qubit of the two-qubit Werner state leads to sudden death of entanglement when the parameter x lies in the range $\frac{1}{3} < x < \frac{1}{2}$. For $x \geq 0.5$, there is an exponential decay of entanglement.

Now treating the first qubit of Werner state with phase noise we have,

$$\rho_{wp} = \begin{pmatrix} \frac{1-x}{4} & 0 & 0 & 0 \\ 0 & \frac{1+x}{4} & \frac{x\gamma}{2} & 0 \\ 0 & \frac{x\gamma}{2} & \frac{1+x}{4} & 0 \\ 0 & 0 & 0 & \frac{1-x}{4} \end{pmatrix}. \tag{24}$$

$$C_{wp} = \frac{1}{2} \text{Max } [0,\, 2x\gamma - (1-x)] \tag{25}$$

It is clear that, $C_{wp} = 0$ when $2x\gamma - (1-x) \leq 0$ or when $t \geq \frac{2}{\tau} \ln \left[\frac{2x}{1-x} \right]$. One can easily see that finite time disentanglement occurs for all values of x lying in the range $\frac{1}{3} < x < 1$. It is illustrated in Fig. 8.

When we allow the first qubit of the Werner state to interact with depolarizing noise we have the following result,

$$\rho_{wd} = \begin{pmatrix} \frac{3+(4p-3)x}{12} & 0 & 0 & 0 \\ 0 & \frac{3+(3-4p)x}{12} & \frac{(3-4p)x}{6} & 0 \\ 0 & \frac{(3-4p)x}{6} & \frac{3+(3-4p)x}{12} & 0 \\ 0 & 0 & 0 & \frac{3+(4p-3)x}{12} \end{pmatrix} \tag{26}$$

It is easy to caluclate the concurrence and is given by,

$$C_{wd} = \frac{1}{6} \text{Max } [0,\, 2(3-4p)x - (3+(4p-3)x)] \tag{27}$$

It is clear that, $C_{wd} = 0$ when

$$2(3-4p)x - (3+(4p-3)x) \leq 0 \Rightarrow p \geq \frac{3x-1}{4x}$$
$$\Rightarrow t \geq \ln \left[\frac{4x}{x+1} \right] \tag{28}$$



FIG. 8: For phase noise acting on a single qubit of the two-qubit state, there is strictly asymptotic decay of entanglement only when the parameter $x = 1$. Though there is a finite time disentanglement for other values of x i.e., $\frac{1}{3} < x < 1$, the decay of entanglement is smooth and not very sudden.

Here, it is not difficult to see that sudden death of entanglement occurs for all values of x in the range $\frac{1}{3} < x \leq 1$. The result is also illustrated with the following graph.



FIG. 9: Occurence of sudden death of entanglement in Werner state when its first qubit alone is subjected to depolarizing noise

In Table I, we have tabulated the time t_d at which disentanglement occurs for chosen parameters of two-qubit isotropic and Werner states under the action of the three noises considered here.

IV. CONCLUSION

In this article, we have shown that a single noise acting on any one of the qubits of an entangled two-qubit state is sufficient for the occurence of entanglement sudden death. We have illustrated this fact by subjecting one of the qubits of a class of two-qubit X states individually to three different kind of noises, amplitude, dephasing and depolarizing. We have worked out the time at which disentanglement occurs for two special states of the X class of states, the two-qubit isotropic and Werner states. Through this work, we have demonstrated that composite noise is not necessary for sudden death of entanglement and entanglement sudden death is a natural phenomena that can happen even when only one part of a composite entangled system is exposed to a noisy environment.

We wish to emphasize here that while the possibility of finite time disentanglement for two qubit X states under the action of *classical dephasing noise* on only one of the qubits is pointed out by Yu and Eberly in [29], the occurence of ESD due to the application of *local quantum noise* on a qubit of two-qubit enangled states has not been given due atention. We hope our work will initiate discussion in this regard.

TABLE I: Disentanglement times for Isotropic and Werner states under different noisy environments

Noise	State	Parameter x	Disentanglement time τt_d	State	Parameter x	Disentanglement time τt_d
Amplitude	Isotropic	0.55	0.405	Werner	0.4	0.405
		0.6	1.38		0.499	4.83
		$\geq$0.625	∞		$\geq$0.5	∞
Phase	Isotropic	0.65	1.653	Werner	0.5	1.386
		0.75	2.773		0.65	2.624
		0.85	4.16		0.85	4.855
		1	∞		1	∞
Depolarising	Isotropic	0.65	0.33	Werner	0.45	0.216
		0.75	0.47		0.65	0.455
		0.85	0.575		0.85	0.61
		1	0.693		1	0.693

Acknowledgement:

Yashodamma K.O acknowledges the support of Department of Science and Technology (DST), Govt. of India in the form of INSPIRE fellowship.

[1] E. Schrödinger, Naturewissenschaften, **23**, 807 (1935)
[2] A. Einstein, B. Podolsky, N. Rosen, Phys. Rev. **47**, 777 (1935)
[3] M. A. Nielsen, I.I. Chuang, Quantum Computation and Quantum Information (Cambridge Univ. Press, Cambridge, 2000)
[4] C. H. Bennett, D.P. DiVincenzo, Nature **404**, 247 (2000)
[5] W. H. Zurek, Rev. Mod. Phys. **75** 715 (2003).
[6] M. Schlosshawer, Rev. Mod. Phys. **76**, 1267 (2005).
[7] C. Simon and J. Kempe, Phys. Rev. A **65** , 052327 (2002).
[8] W. Dür and H. J. Briegel, Phys. Rev. lett. **92**, 180403 (2004).
[9] L. Aolita, R. Chaves, P. Cavakanti, A. Acin and L. Davidovich, Phys. Rev. Lett. **100**, 080501(2008).
[10] T. Yu and J.H. Eberly, Phys. Rev. Lett. **93**, 140404 (2004).
[11] T. Yu and J.H. Eberly, Phys. Rev. Lett. **97**, 140403 (2006).
[12] M. Siomau and Ali. A. Kamli, quant-ph/arxiv:1205.6601.
[13] K. Roszak, P. Machnikowski, Phys. Rev. A 73 (2006) 022313.
[14] M. Yönac, T. Yu and J.H. Eberly, J. Phys. B. **39**, S621, (2006)
[15] T. Yu and J. H. Eberly, Science 316, 555 (2007).
[16] T. Yu and J. H. Eberly, Quantum Inform. and Comput. **7**, 459 (2007)
[17] H.T. Cui, K. Li and X.X. Yi, Phys.Lett.A, **365** (2007).
[18] M.P. Almeida, F. de Melo, M. Hor-Myell, A. Salles, S.P. Walborn, P.H. Souto Ribero, L. Davidovich, Science, **316**,579 (2007).
[19] K. Ann, G. Jaegar, Phys. Rev. A **76**, 044101 (2007)
[20] Xiao-Feng Qian and J.H. Eberly, quant-ph/arxiv:1209.3005.
[21] K. Ann and G. Gaegar, Phys.Lett.A **372**, 579 (2008)
[22] S. Khan, quant-ph/arxiv:1012:1028.
[23] S. Hill and W. K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997).
[24] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
[25] M. Horodecki and P. Horodecki, Phys. Rev. A. **59**, 4206 (1999).
[26] E. Chitambar, arXiv:1110.3057.
[27] R. F. Werner Phys. Rev. A **40**, 4277 (1989).
[28] D. Bruss, J. Math. Phys. **43**, 4237 (2002)
[29] T. Yu and J.H. Eberly, Optics Comm., **264**, 393 (2006).